

GC/MS Analysis of Essential Oil Isolated from the Roots of *Cymbopogon Winterianus* Jowitt

AJAY KUMAR¹ and ABHA SHUKLA²

¹Department of Applied Chemistry,
Faculty of Engineering and Technology,
Gurukul Kangri University,
Haridwar, Uttarakhand, INDIA.

²Department of Chemistry,
Kanya Gurukul Campus, Gurukul Kangri University,
Haridwar, Uttarakhand, INDIA.

(Received on: January 10, 2014)

ABSTRACT

Twelve sesquiterpenoids and six known hydrocarbons were identified in essential oil from the roots of *Cymbopogon winterianus* Jowitt by gas chromatography-mass spectrometry (GC/MS) method, of which the major constituents were α -elemol (32.26%), guaialol (18.81%), methylcyclohexane (9.24%), γ -eudesmol (8.53%), τ -muurolol (6.83%), β -elemene (6.37%), δ -cadinene (4.23%) and (+)-citronellol (2.56%). The essential oil yield and the percentage of identified compounds were 0.16% and 100% respectively. The sesquiterpenoid constituents amounted to 86.58% of the oil. The oil consist several flavor and fragrance compounds and GC/MS findings suggest that the oil may be used in soap, cosmetics, perfumery and food flavoring industries.

Keywords: *Cymbopogon winterianus* Jowitt, Roots, Essential oil, GC/MS, Sesquiterpenoids.

INTRODUCTION

Essential oils are well known for their flavor and fragrance properties. They have great importance due to their versatile applications in soap, food, cosmetics, perfumery and pharmaceutical industries¹.

Some fragrance oils are widely used in aromatherapy to reduce anxiety, stress, and depression of patients², whereas some oils have amazing skin regenerating potential and used in massage after radiotherapy of a cancer patient to reduce its side effects³. Essential oils are composed by highly

bioactive molecules and simply isolated from plant parts by steam distillation or hydro-distillation method. Recent investigations revealed the antibacterial, antifungal, antiviral, anti-inflammatory, anti-tuberculosis, anti-carcinogenic, anticonvulsant, anti-nociceptive, anti-hypertensive, analgesic, anti-oxidant, larvicidal and insecticidal properties of essential oils⁴⁻⁷.

Cymbopogon winterianus Jowitt (Java citronella) is an aromatic plant growing in India, Sri Lanka, Central America, Taiwan, Brazil, Congo, East Africa, Guatemala and West Indies⁸. It is a stemless perennial herb belonging to the Poaceae family of *Cymbopogon* species. The leaves oil of *Cymbopogon winterianus* Jowitt is popular for its aroma and biological properties like as anti-nociceptive, antimicrobial, anticonvulsant and mosquito repellent⁹⁻¹³. In the present study the chemical composition of essential oil isolated from the roots of *Cymbopogon winterianus* Jowitt was investigated by GC/MS method which has not been exploited before.

EXPERIMENTAL

Collection of Plant Material

Fresh roots of *Cymbopogon winterianus* Jowitt were collected from Dr. Shushila Tiwari herbal garden Rishikesh, India in April, 2012. The plant was taxonomically identified and authenticated by the Botanical Survey of India, Dehradun, Uttarakhand by receiving a voucher specimen (Acc. No. 114828).

Isolation Procedure

Fresh roots of plant were washed in tap water and shade dried for 30 days. Dry roots were crushed by using a mechanical blender. 50gm of crushed roots of *Cymbopogon winterianus* Jowitt were subjected to steam distillation for 8 h by using a Clevenger apparatus (Clevenger, 1928). The oil was extracted from the distillate with n-hexane and dried over anhydrous sodium sulfate. The isolated oil was stored in an airtight dark colored bottle with proper labeling for GC/MS analysis.

Gas Chromatography-Mass Spectrometry Analysis

The essential oil from the roots of *Cymbopogon winterianus* Jowitt was analyzed on a Shimadzu QP-2010 (Shimadzu Corporation, Japan) system comprising an autosampler and gas chromatograph coupled to a mass spectrometer (GC/MS) employing the following conditions: Rxi-5 Sil MS cross bond, selectively composed of 5% diphenyl/95% dimethyl polysiloxate capillary column (30 m length x 0.25mm diameter x 0.10 μ m thickness), helium (99.999% purity) was used as carrier gas at a constant flow of 0.7 ml/min and an injection volume of 1.0 μ l of diluted oil in n-hexane was employed (split injection ratio of 1:20), injector temperature 250°C and ion source temperature 200°C. The GC oven temperature was programmed from 50°C (5 min) to 280°C (10 min) at a rate of 4°C/min. The mass spectra were taken at 70 eV (EI mode) with scanning range of m/z 30-600 (interface line temperature 300°C).

Identification of Compounds

The individual constituents of essential oil were identified by computerized matching of their mass spectra and retention indices with those gathered in the NIST 08, FFNS 1.2 and WILEY 8-Mass Spectral libraries of the GC/MS data system.

RESULTS AND DISCUSSION

The volatile oil content obtained

from the roots of Java citronella was 0.16% v/w, and it exhibited yellow color with lemony odor. The results of chemical composition study of essential oil from the roots of *Cymbopogon winterianus* Jowitt are shown in Table 1. The major constituents were α -elemol (32.26%), guaialol (18.81%), methylcyclohexane (9.24%), γ -eudesmol (8.53%), τ -muurolol (6.83%), β -elemene (6.37%), δ -cadinene (4.23%) and (+)-citronellol (2.56%).

Table 1. Chemical composition of essential oil obtained from the roots of *Cymbopogon winterianus* Jowitt

No.	Compound	RT ^a	% Area	MW ^b	MF ^c
1.	Methyl cyclohexane	2.667	9.24	98	C ₇ H ₁₄
2.	Ethyl cyclohexane	2.717	0.77	98	C ₇ H ₁₄
3.	1,2,4-Trimethyl cyclohexane	2.763	0.44	112	C ₈ H ₁₆
4.	Benzene, methyl-	2.983	0.94	92	C ₇ H ₈
5.	1,3-Dimethylcyclohexane	3.137	0.92	112	C ₈ H ₁₆
6.	Nonane	3.360	0.75	128	C ₉ H ₂₀
7.	(+)-Citronellol	16.697	2.56	156	C ₁₀ H ₂₀ O
8.	β -Elemene	21.360	6.37	204	C ₁₅ H ₂₄
9.	E-Germacrene D	23.693	1.34	204	C ₁₅ H ₂₄
10.	β -Selinene	23.890	0.85	204	C ₁₅ H ₂₄
11.	α -Muurolene	24.197	1.37	204	C ₁₅ H ₂₄
12.	γ -Cadinene	24.557	1.84	204	C ₁₅ H ₂₄
13.	δ -Cadinene	24.763	4.23	204	C ₁₅ H ₂₄
14.	α -Elemol	25.713	32.26	222	C ₁₅ H ₂₆ O
15.	cis-Nerolidol	27.130	1.59	222	C ₁₅ H ₂₆ O
16.	γ -Eudesmol	27.670	8.53	222	C ₁₅ H ₂₆ O
17.	τ -Muurolol	27.810	6.83	222	C ₁₅ H ₂₆ O
18.	Guaialol	28.297	18.81	222	C ₁₅ H ₂₆ O
Total Identified Constituents		100.00 %			

^a Retention Index relative to C₅-C₃₂ n-alkanes as standard; Rxi-5 Sil MS column; identification was done by GC/MS library

^b Molecular mass

^c Molecular formula




Figure 1. Structures of identified sesquiterpenes from the essential oil of *Cymbopogon winterianus* Jowitt roots

α -Elemol was found as a major constituent, reported as a floral odorous chemical and make the oil applications in soap and perfumery as fixatives (H. R. Ansari *at el.*, 1974). Guaiol is used in the synthesis of some important chemicals like as guaiazulene which is an anti-inflammatory¹⁵. γ -Eudesmol has been reported as a perfumery chemical with sweet-woody and warm odor^{14,16}. τ -Muurolool has been reported for its antioxidant, anti-cancer and antifungal properties¹⁷⁻¹⁹. β -Elemene is a floral compound with sweet odor and reported as a potential chemopreventive anti-cancer agent²⁰⁻²¹. δ -Cadinene is a natural enzyme which participates in terpenoid biosynthesis²². (+)-Citronellol has been reported as a fragrance and mosquito repellents chemical²³. Nerolidol is an important flavoring and perfumery chemical with a woody floral aroma. It also used as a skin penetration enhancer for the transdermal delivery of drugs²⁴⁻²⁵. E-Germacrene D has a faint, sweet-woody odor which is an important insect pheromone¹⁴.

Results showed the most identified compounds in the *Cymbopogon winterianus* Jowitt roots oil were sesquiterpenes hydrocarbons and accounted to 86.58% of the oil. The structures of identified sesquiterpenes are presented in Figure 1. The identified sesquiterpenes are the main aroma compounds and most of them are biologically active, therefore the oil may be a vital ingredient of soap, candles, cosmetics, perfumes and pharmaceuticals¹⁴.

CONCLUSION

The essential oil from the roots of *Cymbopogon winterianus* Jowitt was

isolated by steam-distillation and analyzed by gas chromatography-mass spectrometry (GC/MS). Results showed that twelve sesquiterpenes which accounts to 86.58% of the oil were isolated from the essential oil of java citronella roots. Sesquiterpenes may be accountable for the biological properties of the oil. The results conclude the identified sesquiterpenes in essential oil studied possess characteristic aroma and several biological active compounds; hence the oil may be used widely in soaps, candles, flavors, fragrances, pharmaceuticals and in aromatherapy.

REFERENCES

1. K. Husnu Can Baser, Gerhard Buchbauer, *Essential Oils: Science, Technology, and Applications*, CRC Press, Taylor and Francis Group, *Boca Raton*, pp. 118, 187, 550, 549, 553, 563, 844, 881 (2010).
2. M. Sugumaran and T. Vetrichelvan. *Aromatherapy: The Power of Scent. Ethnobotanical Leaflets*, 12: 591-98 (2008).
3. Roger M. Smith. *Essential Oil Analysis by Capillary Gas Chromatography and Carbon-13 NMR Spectroscopy. Biochem Soc Trans.*, 11,610 (1983).
4. R.K. Upadhyay, *Essential Oils: Anti-Microbial, Antihelminthic, Antiviral, Anticancer and Anti-Insect Properties. J. Appl. Biosci.*, 36(1): 1-22 (2010)..
5. Amirhossein Sahebkar, Mehrdad Iranshahi. *Biological Activities of Essential Oils from the Genus Ferula Apiaceae. Asian Biomed.*, 4 (6): 835-847 (2010).
6. Kalembe, D.; Kunicka, A. *Antibacterial and Antifungal Properties of Essential Oils. Curr. Med. Chem.*, 10 (10): 813-829 (2003).

7. Manindra Mohan, S Zafar Haider, Harish Chandra Andola, Vijay Kant Purohit. Essential Oils as Green Pesticides: For Sustainable Agriculture. *Res J Pharm Biol Chem Sci.*, 2(4):100-106 (2011).
8. Baby P. Skaria, P.P. Joy, Samuel Mathew, Gracy Mathew, Ancy Joseph and Ragina Joseph, Textbook of Aromatic Plants, *Horticulture Science Series-I*, New Delhi, pp.77 (2007).
9. Wen-Ru Li, Qing-Shan Shi, You-Sheng Ouyang, Yi-Ben Chen, Shun-Shan Duan. Antifungal Effects of Citronella Oil Against *Aspergillus Niger* ATCC 16404. *Appl. Microbiol. Biotechnol.*, 97(16): 7483-7492 (2013).
10. Silva NCC, Fernandes Junior A. Biological Properties of Medicinal Plants: A Review of their Antimicrobial Activity. *J Venom Anim and Toxins*, 16 (3):402-413 (2010).
11. L. J. Quintans-Júnior, T.T. Souza, B.S. Leite, N.M.N. Lessa, L.R. Bonjardim, M.R.V. Santos, P.B. Alves, A.F. Blank, A.R. Antoniolli, Phytochemical Screening and Anticonvulsant Activity of *Cymbopogon winterianus* Jowitt (Poaceae) Leaf Essential Oil in Rodents. *Phytomedicine*, (15) 8: 619-624 (2008).
12. Usawadee Sakulku, Onanong Nuchuchua, Napaporn Uawongyart, Satit Puttipipatkachorn, Apinan Soottitanawat, Uracha Ruktanonchai. Characterization and Mosquito Repellent Activity of Citronella Oil Nanoemulsion. *Int. J. Pharm.*, 372 (1-2):105-111 (2009).
13. Pal Mahesh, Chandrashekar K.. Mosquito Repellent Activity of Piper Betel Linn. *ijpls*, 313-315 (2010).
14. H. R. Ansari, A. J. Curtis, Sesuiterpenes in the Perfumery Industry. *J. Soc. Cosmet. Chem.*, 25: 203-231 (2010).
15. Kurt Bauer, Dorothea Garbe, Horst Surburg, Common Fragrance and Flavor Materials: Preparation, Properties and Uses, London (1997).
16. Timothy Alan Scavone, Loveland, OH, Jianjun Justin Li, West Chester, OH, Marc Adam Flickinger, Oakley, Jonathan Robert Cettis Mason, Compositions for Delivering Perfume to the Skin, US 2013/0252933 A1 (2013).
17. Cheng SS, Wu CL, Chang HT, Kao YT, Chang ST. Antitermitic and Antifungal Activities of Essential Oil of *Calocedrus Formosana* Leaf and its Composition. *J. Chem. Ecol.* 30(10): 1957-67 (2004).
18. Shang-Tzen Chang, David Sheng-Yang Wang, Chi-Lin Wu, Shine-Gwo Shiah, Yueh-Hsiung Kuo, Ching-Jer Chang. Cytotoxicity of Extractives from *Taiwania Cryptomerioides* Heartwood. *Phytochemistry*, 55:227-232 (2000).
19. Sukanya Keawsa-ard and Samart Kongtaweelert. Antioxidant, Antibacterial, Anticancer Activities and Chemical Constituents of the Essential Oil from *Mesua ferrea* Leaves. *Chiang Mai J. Sci.*, 39(3): 455-463 (2012).
20. Edris, Amr E. Anti-Cancer Properties of *Nigella* spp. Essential Oils and their Major Constituents, Thymoquinone and β -Elemene. *Current Clinical Pharmacology*, 4(1): 43-46 (2009).
21. Lei Tao, Liang Zhou, Luying Zheng, Min Yao. Elemene Displays Anti-Cancer Ability on Laryngeal Cancer Cells in Vitro and in Vivo. *Cancer Chemother Pharmacol.*, 58:24-34 (2006).

22. Davis GD, Essenberg M. "(+)-delta-Cadinene is a Product of Sesquiterpene Cyclase Activity in Cotton". *Phytochemistry* 39 (3): 553–567 (1995).
23. Taylor WG, Schreck CE. Chiral-phase Capillary Gas Chromatography and mosquito Repellent Activity of Some Oxazolidine Derivatives of (+)- and (-)-Citronellol. *J. Pharm Sci.*, 74 (5): 534-539 (1985).
24. Byron F. Brehm-Stecher and Eric A. Johnson. Sensitization of *Staphylococcus Aureus* and *Escherichia Coli* to Antibiotics by the Sesquiterpenoids Nerolidol, Farnesol, Bisabolol, and Apitone. *Antimicrob. Agents Chemother*, 47:103357-103360 (2003).
25. Denise C. Arruda, Fabio Luiz D'Alexandri, Alejandro M. Katzin and Silvia R. B. Uliana. Antileishmanial Activity of the Terpene Nerolidol. *Antimicrob. Agents Chemother*, 49 (5): 1679-1687 (2005).